


Multiple Risk Factors Analysis of Post-dural Puncture Headache (PDPH) among Parturient Patients Underwent Caesarean Section at Obstetric Hospitals in Somaliland



Ghazi Mohamed^{1,*} , Khalid Abbas Owish Sukar¹, Ekram Said Hussein Guled², Sara Musa Abdalla Elamin¹, Praveen Kumar Kandakurthi³, Murtada Fathelrahman¹, Mohamed Salah Bashir⁴ and Halah Salah Mohammed Abdalaziz⁵

¹Anesthesia Technology Department, College of Health Sciences, Gulf Medical University, United Arab Emirates

²Anesthesia, Haldor multi-specialty Hospital, Somalia

³Dean College of Health Sciences, Gulf Medical University, United Arab Emirates

⁴King Saud Bin AbdulAziz, University of Health Sciences Anesthesia Technology Program, Saudi Arabia

⁵Anaesthesia Technology Department, College of Applied Sciences, Almaarefa University, Saudi Arabia

Abstract:

Background: Regional anesthesia, particularly spinal technique, is administered during obstetric surgery. The patient and the procedure have risk factors that affect the likelihood of post-dural puncture headaches (PDPH). This study aimed to assess multivariate statistical analyses of risk factors that predict PDPH occurrence in patients who underwent Caesarean section in an obstetrics hospital in Somaliland.

Methods: A cross-sectional study was designed and conducted in obstetrics-selected hospitals in Somaliland. Four hundred parturient patients were included. Statistical tests included Chi Square and risk analysis (Odds & Risk relative ratios) and the Receiver Operating Characteristic (ROC) curve was used to analyze sensitive risk factors to predict PDPH occurrence.

Results: The results revealed that 325 parturient patients (81.25%) developed PDPH while 75 parturient patients (18.75%) didn't report PDPH; multiple lumbar puncture attempts were significantly associated with an increased risk of PDPH, with a P-value of 0.001 and an odds ratio (OR) of 2.705 (95% CI: 1.588–4.605). The ROC analysis for these attempts showed an area under the curve (AUC) of 62%, indicating that the sensitivity and specificity of lumbar puncture attempts (56%, 32%) to predict PDPH. However, other risk factors (Age, PMI, Needle type & size, patient's position) indicate weak predictive value for PDPH occurrence.

Conclusion: The study concluded that Lumbar puncture attempts were a significant risk factor in predicting PDPH compared to other. However, the other risk factors were reported as weak, sensitive and specificity risk factors to predict PDPH occurrence.

Keywords: Complications of spinal anesthesia, Postural puncture headache (PDPH), PDPH risk factors, Caesarean section, Receiver Operating Characteristic, Area under the curve.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Anesthesia Technology Department, College of Health Sciences, Gulf Medical University, United Arab Emirates; E-mail: ghazi@gmu.ac.ae

Cite as: Mohamed G, Sukar K, Guled E, Elamin S, Kandakurthi P, Fathelrahman M, Bashir M, Mohammed Abdalaziz H. Multiple Risk Factors Analysis of Post-dural Puncture Headache (PDPH) among Parturient Patients Underwent Caesarean Section at Obstetric Hospitals in Somaliland. *Open Anesthesiol J*, 2024; 18: e25896458339013. <http://dx.doi.org/10.2174/0125896458339013241009102454>



Received: July 16, 2024
Revised: September 12, 2024
Accepted: October 02, 2024
Published: October 17, 2024



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Cesarean section (CS) under spinal anesthesia has advantages such as allowing the mother to stay conscious throughout the procedure, minimal depression of the fetus, small drug dose, low failure rate, rapid onset, and simplicity of the technique. Therefore, since 2001, performing CS under spinal anesthesia has become a popular and preferred method of general anesthesia [1, 2].

A frequent and uncomfortable complication of spinal anesthesia is post-dural puncture headache (PDPH). It is a throbbing pain on both sides of the head in the front-back area that arises five days after a dural puncture. The pain can get worse with activities that increase pressure in the head, such as sneezing, coughing, pushing, and sudden movements, but resting down can alleviate the pain a bit within 15 minutes [3]. When the needle punctures the dura and causes a hole, cerebrospinal fluid (CSF) can escape rapidly through it, which lowers the pressure inside the skull and leads to post-dural puncture headache (PDPH). Signs of PDPH include feeling dizzy, having a stiff neck, being sensitive to light, seeing double, feeling nauseous, and throwing up [4]. With good rest, fluids, and painkillers, within 48 hours [5], these signs usually diminish in a week. Some factors increase the chance of obstetric patients getting PDPH, such as age, weight, pregnancy, or previous headaches or migraines. The likelihood of PDPH also varies with the needle's type and size, how it is inserted, how many tries it takes, and how much CSF fluid is lost during the procedure [6]. Needles with pencil-point tips (Sprotte or Whitacre) that are 25F to 27F in size produce fewer PDPH than needles with Quincke cutting tips because their tips are blunter and narrower. Thus, the needle size and type are the main determinants of PDPH, and using a smaller needle diameter significantly lowers the PDPH risk [7]. The study aims to determine the sensitivity and specificity of different risk factors for PDPH prediction in obstetric patients with spinal anesthesia. There is not much information on some aspects, such as the number of pregnancies. The incidence and risk factors of PDPH after different neuraxial anesthesia methods are still being researched. Moreover, those studies do not concentrate on specific risk factors' sensitivity, specificity, and risk analysis that can predict the occurrence of PDPH.

Following the administration of regional anesthesia, the patient and the procedure tend to have risk factors that affect the likelihood of post-dural puncture headaches, particularly spinal anesthesia. Factors that affect the risk represent the needle's characteristics and how often the lumbar puncture is attempted, age, BMI, and headache history, including post-dural puncture headache. The goal of the study was to ascertain, which had not been documented in prior studies, the sensitivity-specificity risk factors analysis that can predict the incidence of PDPH. One study found that PDPH can result from different spinal surgeries and neuraxial anesthesia combinations, and some risk factors are known. How often these risk factors affect PDPH after various treatments is not well-studied [8]. Thus, the study identified the main

risk factors in multiple analyses to forecast the PDPH incidence that are accurate and precise, using a method based on scientific evidence that could lower PDPH in the future.

1.1. Objectives

- To evaluate the risk factors of PDPH by using multivariate risk statistical analyses such as risk and odd ratio sensitivity and specificity by ROC curve among caesarean section parturients in Somaliland.

2. METHOD

2.1. Study Design and Participants

In this cross-sectional study, 400 parturient patients underwent cesarean section and were anesthetized by spinal anesthesia at Hargeisa Group Hospital, Gargaar Multi-specialty, and Kaah Community Hospitals from July 2023 to June 2024. Parturient patients aged between 17 to 45 years were included and were categorized according to the activity of reproduction to group age [(17- 29 years) with a peak of fertility, group (30-35 years) decreased fertility, and group (greater than 35 years) as declined fertility and reproduction] [9, 18]. Maternal patients ASA class I - II were planned for scheduled caesarean sections during the period of study.

2.2. Ethical Considerations

This study was approved by the Department of Planning/Policy and Strategic Information, National Ethical Committee, along with a unit of research development, ministry of Health Development, Republic Somaliland (Project ref: MOHD/DG:2/1246/2023).

2.3. Setting and Study Size

400 Parturient patients underwent cesarean section and were anesthetized by spinal anesthesia at Hargeisa Group Hospital, Gargaar Multi-specialty, and Kaah Community Hospitals, which were conducted from July 2023 to June 2024. Patients admitted to the hospital for spinal anesthesia during the data collection period were included in the study. The sample size was calculated based on the formula for unmatched case-control studies.

Formula for Sample Calculation: $N = Z^2 Pq / e^2$

N = sample size

Z = Number of patients admitted for LSCS under spinal anesthesia
Pq = Variance of population

E = Allowable error 5%

N= Maternal patients

2.4. Technique of Spinal Anesthesia

Spinal anesthesia was performed by injecting 12-15 mg of Bupivacaine 0.5% isotonic at L3-L4 according to the weight. The spinal needle size ranging from 25, 26, and 27 G was used during anesthesia and most of the patients were given anesthesia in a sitting position respective to the lateral. Two types of spinal needles were used for the lumbar puncture technique, including pencil point and

quincke tip and patient vital signs were monitored by standard anesthesia monitors during the intra-operative and postoperative periods.

2.5. Variables, Data Sources/measurement, and Bias

Demographic data and consent regarding parturient patients were collected preoperatively in antenatal wards through parameters of risk factors of post-dural puncture headache (spinal needle type, size, and attempt of lumbar puncture intra-operatively) from anesthesia records and anesthesia team and postoperatively after 24 to 3 days by stranded questionnaire from the patients in postnatal wards.

This data was collected by a team of trained anesthesia practitioners in the hospitals included in this study. Data for this research was acquired through a questionnaire and is stored safely in an online database in an Excel spreadsheet available upon request.

Data was collected equally and confidentially from the hospital selected in the study.

2.6. Quantitative Variables

This study included 400 parturient patients who underwent cesarean section under spinal anesthesia and the study reported that 325 (81.25%) patients developed PDPH and 75 (18.75%) patients did not (Fig. 1).

2.7. Statistical Methods

SPSS program version 26.0 was used for statistical analysis. Nominal variables are presented as frequencies and Chi square tests were used for the comparison of

categorical variables between groups. P value \leq (0.05) was expressed as significance level, and risk analysis of risk factors by Odds ratio & Relative risk ratio with 95% CI was measured and sensitivity and specificity estimated by Receiver Operating Characteristics (ROC) curve analysis cut-off value area under the curve of results parameter. The results were considered positive if there were less or equal cut-offs or negative when there were more cut-off values and results were presented in tables and figures.

3. RESULTS

3.1. Participants

In this study, 400 patients were included who underwent cesarean section under spinal anesthesia. It was found that 325 (81.25%) patients developed PDPH and 75 (18.75%) patients did not (Fig. 1).

3.2. Descriptive Data

The patient risk factors were grouped into two age groups according to fertility: group age 17- 29 years with peak fertility and group age 30- 35 years with decreased and declined fertility. The age group increased the risk of PDPH by 1.377 times, body mass index increased the PDPH (normal and obese) by 1.753 times, needle type by 1.628 times, and needle size (25G or larger) increased the risk of PDPH by 1.409 times. Patient position (sitting vs. lateral) reduced the risk of PDPH by 0.451 times. Additionally, lumbar puncture attempts were associated with a 2.705 times higher risk of PDPH, as shown in Table 1.




Fig. (1). Shows the postdural puncture headache occurrence among parturient patients in obstetric hospitals in somaliland.

Table 1. Shows the odds ratios of risk factors of post-dural puncture headache (PDPH) among caesarean section patients.

Risk Factors of PDPH	Odds Ratio	95% Confidence Interval	
		Lower	Upper
Age group (17 - 29 & 30-45 years)	1.377	0.824	2.300
Body Mass Index (BMI) (Normal weight / Obese)	1.753	0.334	9.216
Spinal Needle type (Quincke tip / Pencil tip)	1.268	0.472	3.411
Needle size (25 G / > 25 G)	1.409	0.780	2.545
Position (Sitting & Lateral Position)	0.451	0.110	1.848
Lumbar Puncture attempt (1st attempt / More attempts)	2.705	1.588	4.605

Table 2A. Shows relative risk ratios of risk factors (Patient Factors) of postdural puncture headache (PDPH) among caesarean section patients.

Risk Factors of PDPH (Patient Factors)	PDPH Occurrence		P-value	Relative Risk Ratio	95% Confidence Interval	
	No (%)	Yes (%)			Lower	Upper
Age group (17-29 Yrs)	46 (20.9)	174(79.1)	0.24	1.146	0.932	1.408
Age group (30-35years & >35 years)	29 (16.1)	151(83.9)		0.832	0.612	1.132
Body Mass Index (BMI) Normal weight	2(28.6)	5(71.4)	0.50	1.733	0.343	8.763
Body Mass Index (BMI) Obese	73(18.6)	320(81.4)		0.989	0.950	1.029
Setting Position	72 (20.3)	319 (79.7)	0.25	0.978	0.932	1.027
Lateral Position	3 (33.3)	6 (66.7)		2.167	0.554	8.467

Note: Keys: P value ≥ 0.05 means (significant level).

Table 2B. Shows relative risk ratios of risk factors (Technique Factors) of postdural puncture headache (PDPH) among caesarean section patients.

Risk Factors of PDPH (Technique Factors)	PDPH Occurrence		P-value	Relative Risk Ratio	95% Confidence Interval	
	No (%)	Yes (%)			Lower	Upper
Spinal needle (Quincke tip)	70 (19.0)	298 (81.0)	0.63	1.217	0.530	2.798
Spinal needle (Pencil tip)	5(15.6)	27(84.4)		0.960	0.820	1.123
Spinal Needle Size (25 G)	58 (77.3)	230(70.8)	0.25	1.093	0.949	1.258
Spinal Needle Size (> 25 G)	17(22.7)	95(29.2)		0.775	0.494	1.217
Lumbar Puncture (1st attempt)	51(68.0)	143(44.0)	0.001	1.545	1.268	1.884
Lumbar Puncture (Multiple attempts)	24 (32.0)	182 (56.0)		0.571	0.405	0.806

Note: Keys: P value ≥ 0.001 means (significant level).

3.3. Outcome Data, Main Results, and Other Analyses

The results of the current study revealed that there were no significant changes (P value 0.24) of PDPH occurrence between age groups; however, the age group 17-29 years reported a greater risk ratio of PDPH (1.146) compared to age group 30-45 year, which reported 0.832, BMI also reported no significant changes of PDPH for normal and obese patients (P value 0.50, RR 1.733 & 0.989) and both setting and lateral positions were detected to have no significant changes (P vale 0.25) and slightly increased of relative risk ratio in lateral position 2.167 respective to other position have shown in Table 2A.

The results of the current study revealed that there were no significant changes (P value 0.63) in PDPH occurrence related to needle types; however, quincke tip

spinal needle statistically reported slight increases in the PDPH occurrence and risk ratio (1.217) compared to pencil tip (0.960). Spinal needle size 25G reported slight statistical increases in the PDPH occurrence and risk ratio (81% RR ratio 1.217) compared to other spinal needle sizes but there was a significant association. However, lumber puncture attempts reported a significant elevation of PDPH occurrence with multiple attempts compared to the first attempt have shown in Table 2B.

The result variable(s): Age, Body Mass Index (BMI), and position have shown area under the curve. Age group 53% (weak), sensitivity 45%, specificity 38%, ROC curve cut off for the BMI is 52% (weak), the result area under a curve of Position 0.48 (48%) (weak) as shown in Fig. (2).

3.4. Study Design and Participants

Regarding the ROC curve, the cut-off value of LP attempts (AUC, the area under the curve is 62%) sensitivity of 56% and specificity of 32%. LP attempts result is considered positive if it is less or equal cut-off or

negative when there are more cut-off values. The ROC curve cut-off value area under the curve of needle type and needle size is 50% and 53% (Weak). The results of needle type and size are considered positive if it is less or equals the cut-off or negative when there are more cut-off values shown in Fig. (3).




Fig. (2). Receiver operating characteristics (ROC) of the age, body mass index (BMI), and position of parturient patients.




Fig. (3). Receiver operating characteristics (ROC) of the needle type, needle size, and lumbar puncture attempts of parturient patients.

4. DISCUSSION

PDPH is a common complication of neuraxial anesthesia that develops primarily after puncturing the dura mater, causing a CSF leakage, which results in intracranial hypotension [10]. Multiple reports of research documented the risk factors that influence PDPH occurrence, including patient's risk factors (age, BMI, ASA classification, gestational age, urgency of surgery, previous PDPH, parity, and gravidity) [11] and technique's risk factors (needle type, needle gauge, number of attempts, and direction of bevel) [12].

4.1. Key Results

The purpose of this study, which involved 400 parturient patients who underwent caesarean sections, was to assess the multivariate statistical risk analysis to predict highly sensitive and specific risk factors in developing post-dural puncture headache (PDPH) following spinal anesthesia. Ljubisavljevic *et al.*, and Al-Hashel *et al.*, reported that younger patients had a higher chance of developing PDPH compared to older patients; the possible cause could be related to less CSF production and elastic dura matter [13, 14] and these findings agree with our results which reported that patients between 17-29 years have higher PDPH occurrence when compared with other age groups.

The current results did not detect a significant association between BMI and PDPH occurrence in parturient patients. However, many studies reported that a BMI greater than 30 kg/m² had a significant association with the incidence of PDPH [11, 13, 15].

This study reported that there was no significant association between patient positions during lumbar puncture, needle types and size relevant to PDPH occurrence; however, the relative risk factors statistically were detected between lateral and setting position and different needle sizes to increase risk of PDPH occurrence. The decrease of spinal size to less than 27G for cesarean section reduced PDPH incidence [16]. Moreover, the previous randomized trial reported the incidence of PDPH with needle-type Quincke was 22.43% [17].

The current study detected that spinal needle size 25G reported slight statistical increases in the PDPH occurrence and risk ratio (RR: 1.093; 95% CI =0.949-1.258) compared to other sizes. A smaller needle size reduces the risk of CSF leakage and PDPH when piercing the dura mater [19, 20]. A 27G is better than 25G, as two studies showed less PDPH with 27G ($p < 0.05$) [21, 22]. A similar study also showed that PDPH was more common with 20G and 22G needles than with 24G and 25G (AOR = 4.206; 95% CI = 1.247-14.187; $p = 0.021$), which matches our findings [23].

The findings of this study confirm that the technique-related risk factors, lumbar puncture attempts reported significant increases in PDPH occurrence with both first and multiple attempts and (sensitivity 56% specificity 32% and significant P value 0.001, OR 2.705 95% IC 1.588-4.605) increased incidence and risk of PDPH

occurrence. However, higher incidence reported during multiple attempts in the current results. Every time the dura matter is penetrated, there is more CSF loss and a greater risk of CSF hypotension and PDPH [24]. This is consistent with multiple studies that associate the frequency of dura matter punctures with the occurrence of PDPH, such as Wegi Girma *et al.*, who report that PDPH occurrence increases 1.5 times with multiple attempts ($P = 0.001$) [11]. Another statement from Demilew BC *et al.* is that the chance of PDPH increases 4.7 times with repeated punctures of the dura matter ($P = 0.05$).

CONCLUSION

This study concluded that multiple analyses reflected that Lumbar puncture attempts, particularly multiple attempts, were significant potential risk factors that can predict post-dural puncture headache. However, other factors detected weak sensitivity and specificity.

LIMITATIONS

The study's limitations are that awareness of preoperative consent is not applicable in most hospitals in Somaliland as a mandatory protocol in theater. The limited appropriate quality of technique.

WHAT IS ALREADY KNOWN ABOUT THIS TOPIC

PDPH is a frequent and uncomfortable complication of spinal anesthesia, characterized by throbbing pain on both sides of the head.

Risk Factors: Factors such as age, gender, weight, pregnancy, previous headaches, needle type and size, insertion technique, and number of lumbar puncture attempts influence the likelihood of PDPH.

WHAT THIS STUDY ADDS

High Incidence of PDPH: 81.25% of the participants developed post-dural puncture headache (PDPH) after undergoing a Caesarean section with spinal anesthesia.

Significant Risk Factor: Multiple lumbar punctures attempt significantly increased the likelihood of PDPH occurrence.

Needle Characteristics: The type and size of the needle used in spinal anesthesia play a crucial role in the risk of developing PDPH.

Predictive Analysis: The study used statistical methods like Chi-square, odds ratios, and ROC curves to analyze and predict PDPH risk factors.

AUTHOR'S CONTRIBUTION

H. S., M. A., M. S. B., M. F., P. K. K. and G. M.: Wrote the paper; K. A. and O. S.: Data analysis or interpretation were contributed; E. S. H. G.: Collected the data; S. M. A. E.: Provided the study concept and design.

LIST OF ABBREVIATIONS

PDPH	=	Post-dural Puncture Headache
ROC	=	Receiver Operating Characteristics
BMI	=	Body Mass Index

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Department of Planning/Policy and Strategic Information, National Ethical Committee, along with a unit of research development, ministry of Health Development, Republic Somaliland (Project ref: MOHD/DG:2/1246/2023).

HUMAN AND ANIMAL RIGHTS

All procedures performed in studies involving human participants were in accordance with the ethical standards of institutional and/or research committees and with the 1975 Declaration of Helsinki, as revised in 2013.

CONSENT FOR PUBLICATION

Demographic data and consent regarding parturient patients were collected preoperatively in antenatal wards.

STANDARDS OF REPORTING

STROBE guidelines were followed.

AVAILABILITY OF DATA AND MATERIALS

The data supporting the findings of the article is available in the [Zenodo Repository] at [https://zenodo.org/records/13923127], reference number [10.2174/0125896458339013241009102454, 2024, 18, e25896458339013].

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Idris IM, Weldegiorgis GG, Tesfamariam EH. Maternal satisfaction and its associated factors towards spinal anesthesia for Caesarean Section: A cross-sectional study in two Eritrean Hospitals. *Anesthesiol Res Pract* 2020; 20(2): 1-8. <http://dx.doi.org/10.1155/2020/5025309> PMID: 32256568
- [2] Crawford-Sykes A, Scarlett M, Hambleton IR, Nelson M, Rattray C. Anaesthesia for operative deliveries at the University Hospital of the West Indies: a change of practice. *West Indian Med J* 2005; 54(3): 187-91. <http://dx.doi.org/10.1590/S0043-31442005000300006> PMID: 16209224
- [3] Al-Hashel J, Rady A, Massoud F, Ismail II. Post-dural puncture headache: A prospective study on incidence, risk factors, and clinical characterization of 285 consecutive procedures. *BMC Neurol* 2022; 22(1): 261. <http://dx.doi.org/10.1186/s12883-022-02785-0> PMID: 35836140
- [4] Uppal V, Russell R, Sondokoppam R, *et al.* Consensus practice guidelines on postdural puncture headache from a multisociety, international working group. *JAMA Netw Open* 2023; 6(8): e2325387. <http://dx.doi.org/10.1001/jamanetworkopen.2023.25387> PMID: 37581893
- [5] Chekol B, Yetneberk T, Teshome D. Prevalence and associated factors of post dural puncture headache among parturients who underwent cesarean section with spinal anesthesia: A systemic review and meta-analysis, 2021. *Ann Med Surg* 2021; 66: 102456. <http://dx.doi.org/10.1016/j.amsu.2021.102456>
- [6] Xu H, Liu Y, Song W, *et al.* Comparison of cutting and pencil-point spinal needle in spinal anesthesia regarding postdural puncture headache. *Medicine* 2017; 96(14): e6527. <http://dx.doi.org/10.1097/MD.00000000000006527> PMID: 28383416
- [7] Zorrilla-Vaca A, Mathur V, Wu CL, Grant MC. The impact of spinal needle selection on postdural puncture headache. *Reg Anesth Pain Med* 2018; 43(5): 1. <http://dx.doi.org/10.1097/AAP.0000000000000775> PMID: 29659437
- [8] Jensen RE, Martins N, Parks MM. Public perception of female fertility: Initial fertility, peak fertility, and age-related infertility among U.S. Adults. *Arch Sex Behav* 2018; 47(5): 1507-16. <http://dx.doi.org/10.1007/s10508-018-1197-4> PMID: 29582267
- [9] Ljubisavljevic S. Postdural puncture headache as a complication of lumbar puncture: Clinical manifestations, pathophysiology, and treatment. *Neurol Sci* 2020; 41(12): 3563-8. <http://dx.doi.org/10.1007/s10072-020-04757-z> PMID: 32997283
- [10] Kim M, Yoon H. Comparison of post-dural puncture headache and low back pain between 23 and 25 gauge Quincke spinal needles in patients over 60 years: Randomized, double-blind controlled trial. *Int J Nurs Stud* 2011; 48(11): 1315-22. <http://dx.doi.org/10.1016/j.ijnurstu.2011.04.005> PMID: 21561619
- [11] Weji BG, Obsa MS, Melese KG, Azeze GA. Incidence and risk factors of postdural puncture headache: Prospective cohort study design. *Perioper Med* 2020; 9(1): 32. <http://dx.doi.org/10.1186/s13741-020-00164-2> PMID: 33292510
- [12] Arevalo-Rodriguez I, Muñoz L, Godoy-Casasbuenas N, *et al.* Needle gauge and tip designs for preventing post-dural puncture headache (PDPH). *Cochrane Libr* 2017; 2017(12): CD010807. <http://dx.doi.org/10.1002/14651858.CD010807.pub2> PMID: 28388808
- [13] Zhao G, Song G, Liu J. Efficacy of pharmacological therapies for preventing post-dural puncture headaches in obstetric patients: A Bayesian network meta-analysis of randomized controlled trials. *BMC Pregnancy Childbirth* 2023; 23(1): 215. <http://dx.doi.org/10.1186/s12884-023-05531-7> PMID: 36991366
- [14] Vallejo MC, Zakowski MI. Post-dural puncture headache diagnosis and management. *Baillieres Best Pract Res Clin Anaesthesiol* 2022; 36(1): 179-89. <http://dx.doi.org/10.1016/j.bpa.2022.01.002> PMID: 35659954
- [15] Thakur S, Sharma A, Kaushal S, Sharma A, Sharma N, Thakur PS. Incidence and risk factors of "postdural puncture headache" in women undergoing cesarean delivery under spinal anesthesia with 26g quincke spinal needle, experience of medical college in rural settings in India 2019: A prospective cohort study design. *J Pharm Bioallied Sci* 2022; 14(5) (Suppl. 1): S209-13. http://dx.doi.org/10.4103/jpbs.jpbs_72_22 PMID: 36110769
- [16] Shaikh JM, Memon A, Memon MA, Khan M. Post dural puncture headache after spinal anaesthesia for caesarean section: A comparison of 25 g Quincke, 27 g Quincke and 27 g Whitacre spinal needles. *J Ayub Med Coll Abbottabad* 2008; 20(3): 10-3. PMID: 19610505
- [17] Lakhe G, Shrestha P, Duwadi V. Postdural puncture headache after spinal anaesthesia in parturients undergoing cesarean section in the department of anesthesia in a tertiary care center: A descriptive cross-sectional study. *JNMA J Nepal Med Assoc* 2023; 61(261): 417-20. <http://dx.doi.org/10.31729/jnma.8140> PMID: 37203905
- [18] Li H, Wang Y, Oprea AD, Li J. Postdural puncture headache—risks and current treatment. *Curr Pain Headache Rep* 2022; 26(6): 441-52. <http://dx.doi.org/10.1007/s11916-022-01041-x>
- [19] Doelakeh ES, Chandak A. Risk factors in administering spinal anesthesia: A comprehensive review. *Cureus* 2023; 15(12): e49886. <http://dx.doi.org/10.7759/cureus.49886> PMID: 38174200

- [20] Van de Velde M, Schepers R, Berends N, Vandermeersch E, De Buck F. Ten years of experience with accidental dural puncture and post-dural puncture headache in a tertiary obstetric anaesthesia department. *Int J Obstet Anesth* 2008; 17(4): 329-35. <http://dx.doi.org/10.1016/j.ijoa.2007.04.009> PMID: 18691871
- [21] Traynor AJ, Aragon M, Ghosh D, *et al.* Obstetric anesthesia workforce survey: A 30-year update. *Anesth Analg* 2016; 122(6): 1939-46. <http://dx.doi.org/10.1213/ANE.0000000000001204> PMID: 27088993
- [22] Batova R, Georgiev S. Impact of spinal needle design and approach to postdural puncture headache and spinal anesthesia failure in obstetrics. *Anaesthesiol Intensive Ther* 2019; 51: 77-82. <http://dx.doi.org/10.5114/ait.2019.86166>
- [23] Demilew BC, Tesfaw A, Tefera A, Getnet B, Essa K, Aemro A. Incidence and associated factors of postdural puncture headache for parturients who underwent cesarean section with spinal anesthesia at Debre Tabor General Hospital, Ethiopia; 2019. *SAGE Open Med* 2021; 9 <http://dx.doi.org/10.1177/205031212111051926> PMID: 34676076
- [24] Ali HM, Mohamed MY, Ahmed YM. Postdural puncture headache after spinal anesthesia in cesarean section: Experience in six months in 2736 patients in Kasr El aini teaching hospital - Cairo University. *Egypt J Anaesth* 2014; 30(4): 383-6. <http://dx.doi.org/10.1016/j.egja.2014.06.001>